

ischemia. The demands placed on quality in non-invasive techniques will therefore increase for both echocardiography and nuclear cardiology laboratories.

References

- Laskar R, Grayburn PA: Assessment of myocardial perfusion with contrast echocardiography at rest and with stress: An emerging technology. *Prog Cardiovasc Dis* 43:245-258, 2000
- Tennant R, Wiggers C: The effect of coronary occlusion on myocardial contraction. *Am J Physiol* 112: 351-361, 1935
- Kerber RE, Marcus ML, Ehrhardt J, et al: Correlation between echocardiographically demonstrated segmental dyskinesis and regional myocardial perfusion. *Circulation* 52:1097-1104, 1975
- Sugishita Y, Koseki S, Matsuda M, et al: Dissociation between regional myocardial dysfunction and ECG changes during myocardial ischemia induced by exercise in patients with angina pectoris. *Am Heart J* 106:1-8, 1983
- Corya BC, Rasmussen S, Knoebel SB, et al: M-mode echocardiography in evaluating left ventricular function and surgical risk in patients with coronary artery disease. *Chest* 72:181-185, 1977
- Mason SJ, Weiss JL, Weisfeldt ML, et al: Exercise echocardiography: detection of wall motion abnormalities during ischemia. *Circulation* 59:50-59, 1979
- Battler A, Froelicher VF, Gallagher KP, et al: Dissociation between regional myocardial dysfunction and ECG changes during ischemia in the conscious dog. *Circulation* 62:735-744, 1980
- Sugishita Y, Koseki S: Dynamic exercise echocardiography. *Circulation* 60:743-752, 1979
- Bartel T, Muller S, Borges AC, et al: Left and right heart doppler stress echo in congestive heart failure. *Int J Card Imaging* 10:289-297, 1994
- Parthenakis FI: Assessment of left ventricular ejection dynamics in patients with coronary artery disease during dipyridamole-stress doppler echocardiography. [in process citation]. *Coron Artery Dis* 10:471-477, 1999
- Hunziker PR, Picard MH, Jander N, et al: Regional wall motion assessment in stress echocardiography by tissue doppler bull's-eyes. *J Am Soc Echocardiogr* 12:196-202, 1999
- Leischik R, Bartel T, Mohlenkamp S, et al: Stress echocardiography: New techniques. *Eur Heart J* 18: D49-D56, 1997 (suppl D)
- Katz WE, Gulati VK, Mahler CM, et al: Quantitative evaluation of the segmental left ventricular response to dobutamine stress by tissue doppler echocardiography. *Am J Cardiol* 79:1036-1042, 1997
- Pasquet A, Armstrong G, Beachler L, et al: Use of segmental tissue doppler velocity to quantitate exercise echocardiography. *J Am Soc Echocardiogr* 12: 901-912, 1999.
- Pasquet A, Yamada E, Armstrong G, et al: Influence of dobutamine or exercise stress on the results of pulsed-wave doppler assessment of myocardial velocity. *Am Heart J* 136:753-758, 1999
- Panza JA, Laurienzo JM, Curriel RV, et al: Transesophageal dobutamine stress echocardiography for evaluation of patients with coronary artery disease. *J Am Coll Cardiol* 24:1260-1267, 1994
- Burduli NM, Buziashvili lu I, Kharitonova NI, et al: [stress echocardiography in localization of coronary atherosclerosis and assessment of it severity] [Stress-EkhoKG v opredelenii lokalizatsii i stepeni vyrazhennosti koronarnogo aterosklerozha.]. *Klin Med (Mosk)* 76:22-24, 1998
- Radvan J, Marwick TH, Williams MJ, et al: Evaluation of the extent and timing of the coronary hyperemic response to dipyridamole: A study with transesophageal echocardiography and positron emission tomography with oxygen 15 water. *J Am Soc Echocardiogr* 8:864-873, 1995
- Panza JA, Dilsizian V, Laurienzo JM, et al: Relation between thallium uptake and contractile response to dobutamine. Implications regarding myocardial viability in patients with chronic coronary artery disease and left ventricular dysfunction. *Circulation* 91:990-998, 1995
- Zwas DR, Takuma S, Mullis-Jansson S, et al: Feasibility of real-time 3-dimensional treadmill stress echocardiography. *J Am Soc Echocardiogr* 12:285-289, 1999
- Gibbons EF, Hogan RD, Franklin TD, et al: The natural history of regional dysfunction in a canine preparation of chronic infarction. *Circulation* 71:394-402, 1985
- Meltzer RS, Woythaler JN, Buda AJ, et al: Non-invasive quantification of experimental canine myocardial infarct size using two-dimensional echocardiography. *Eur J Cardiol* 11:215-225, 1980
- Wyatt HL, Meerbaum S, Heng MK, et al: Experimental evaluation of the extent of myocardial dyssynergy and infarct size by two-dimensional echocardiography. *Circulation* 63:607-614, 1981
- Nieminen M, Parisi AF, O'Boyle JE, et al: Serial evaluation of myocardial thickening and thinning in acute experimental infarction: Identification and quantification using two-dimensional echocardiography. *Circulation* 66:174-180, 1982
- Pandian NG, Koyanagi S, Skorton DJ, et al: Relations between 2-dimensional echocardiographic wall thickening abnormalities, myocardial infarct size and coronary risk area in normal and hypertrophied myocardium in dogs. *Am J Cardiol* 52:1318-1325, 1983
- Skorton DJ, Melton HE Jr, Pandian NG, et al: Detection of acute myocardial infarction in closed-chest dogs by analysis of regional two-dimensional echo-

- cardiographic gray-level distributions. *Circ Res* 52:36-44, 1983
27. Pandian NG, Kieso RA, Kerber RE: Two-dimensional echocardiography in experimental coronary stenosis. II. Relationship between systolic wall thinning and regional myocardial perfusion in severe coronary stenosis. *Circulation* 66:603-611, 1982
 28. Heger JJ, Weyman AE, Wann LS, et al: Cross-sectional echocardiography in acute myocardial infarction: Detection and localization of regional left ventricular asynergy. *Circulation* 60:531-538, 1979
 29. Parisi AF, Moynihan PF, Folland ED, et al: Quantitative detection of regional left ventricular contraction abnormalities by two-dimensional echocardiography. II. Accuracy in coronary artery disease. *Circulation* 63:761-767, 1981
 30. Heger JJ, Weyman AE, Wann LS, et al: Cross-sectional echocardiographic analysis of the extent of left ventricular asynergy in acute myocardial infarction. *Circulation* 61:1113-1118, 1980
 31. Horowitz RS, Morganroth J, Parrotto C, et al: Immediate diagnosis of acute myocardial infarction by two-dimensional echocardiography. *Circulation* 65:323-329, 1982
 32. Schiller NB, Shah PM, Crawford M, et al: Recommendations for quantitation of the left ventricle by two-dimensional echocardiography. American society of echocardiography committee on standards, subcommittee on quantitation of two-dimensional echocardiograms. *J Am Soc Echocardiogr* 2:358-367, 1989
 33. Hausnerova E, Gottdiener J, Hecht G, et al: Increased diastolic left ventricular wall thickness during dobutamine stress echocardiography: Effect of afterload. *Circulation* 90:391, 1994 (abstr)
 34. Carstensen S, Ali SM, Stensgaard-Hansen FV, et al: Dobutamine-atropine stress echocardiography in asymptomatic healthy individuals. The relativity of stress-induced hyperkinesia. *Circulation* 92:3453-3463, 1995
 35. Sechtem U, Baer FM, Voth E, et al: Stress functional MRI: Detection of ischemic heart disease and myocardial viability. *J Magn Reson Imaging* 10:667-675, 1999
 36. Scott CH, Sutton MS, Gusani N, et al: Effect of dobutamine on regional left ventricular function measured by tagged magnetic resonance imaging in normal subjects. *Am J Cardiol* 83:412-417, 1999
 37. Mitamura H, Ogawa S, Hori S, et al: Two-dimensional echocardiographic analysis of wall motion abnormalities during handgrip exercise in patients with coronary artery disease. *Am J Cardiol* 48:711-719, 1981
 38. Whitfield S, Aurigemma G, Pape L, et al: Two-dimensional doppler echocardiographic correlation of dipyridamole-thallium stress testing with isometric handgrip. *Am Heart J* 121:1367-1373, 1991
 39. Tawa CB, Baker WB, Kleiman NS, et al: Comparison of adenosine echocardiography, with and without isometric handgrip, to exercise echocardiography in the detection of ischemia in patients with coronary artery disease. *J Am Soc Echocardiogr* 9:33-43, 1996
 40. Picano E, Lattanzi F, Masini M, et al: Does the combination with handgrip increase the sensitivity of dipyridamole-echocardiography test? *Clin Cardiol* 10:37-39, 1987
 41. Cohn PF, Gorlin R, Herman MV, et al: Relation between contractile reserve and prognosis in patients with coronary artery disease and a depressed ejection fraction. *Circulation* 51:414-420, 1975
 42. Helfant RH, De Villa MA, Banka VS: Evaluation of left ventricular performance in coronary heart disease: Use of isometric handgrip stress test. *Catheter Cardiovasc Diagn* 2:59-67, 1976
 43. Mathias W Jr: Safety of dobutamine-atropine stress echocardiography: A prospective experience of 4033 consecutive studies [in process citation]. *J Am Soc Echocardiogr* 12:785-791, 1999
 44. Elhendy A, van Domburg RT, Poldermans D, et al: Safety and feasibility of dobutamine-atropine stress echocardiography for the diagnosis of coronary artery disease in diabetic patients unable to perform an exercise stress test. *Diabetes Care* 21:1797-1802, 1998
 45. Rowlands DB, Ireland MA, Glover DR: The relationship between ambulatory blood pressure and echocardiographically assessed left ventricular hypertrophy. *Clin Sci* 61:101s-103s, 1981 (suppl)
 46. Cusick DA, Bonow RO, Chaudhry FA: Safety of dobutamine stress echocardiography in patients with left ventricular apical thrombus. *Am J Cardiol* 80:1252-1254, 1997
 47. Pellikka PA, Roger VL, Oh JK, et al: Safety of performing dobutamine stress echocardiography in patients with abdominal aortic aneurysm \geq 4 cm in diameter. *Am J Cardiol* 77:413-416, 1996
 48. Hobday TJ, Pellikka PA, Jost CH, et al: Chronotropic response, safety, and accuracy of dobutamine stress echocardiography in patients with atrial fibrillation and known or suspected coronary artery disease. *Am J Cardiol* 82:1425-1427, 1998
 49. Bach DS, Cohen JL, Fioretti PM, et al: Safety and efficacy of closed-loop arbutamine stress echocardiography for detection of coronary artery disease. international arbutamine study group. *Am J Cardiol* 81:32-35, 1998
 50. Cohen JL, Chan KL, Jaarsma W, et al: Arbutamine echocardiography: Efficacy and safety of a new pharmacologic stress agent to induce myocardial ischemia and detect coronary artery disease. The international arbutamine study group. *J Am Coll Cardiol* 26:1168-1175, 1995
 51. Bach DS, Armstrong WF: Adequacy of low-stress arbutamine to provoke myocardial ischemia during echocardiography. international arbutamine study group. *Am J Cardiol* 76:259-262, 1995

52. Picano E, Sicari R, Varga A: Dipyridamole stress echocardiography. *Cardiol Clin* 17:481-499, 1999
53. Picano E, Sicari R, Varga A: Dipyridamole stress echocardiography, in Aurigemma GP (ed): *Stress Echocardiography*. Philadelphia, PA, Saunders, 1999, pp 481-499.
54. Bonow RO, Kent KM, Rosing DR, et al: Exercise-induced ischemia in mildly symptomatic patients with coronary-artery disease and preserved left ventricular function. Identification of subgroups at risk of death during medical therapy. *N Engl J Med* 311:1339-1345, 1984
55. Bach DS: Stress echocardiography for evaluation of hemodynamics: Valvular heart disease, prosthetic valve function, and pulmonary hypertension. *Prog Cardiovasc Dis* 39:543-554, 1997
56. Decena BF 3rd, Tischler MD: Stress echocardiography in valvular heart disease. *Cardiol Clin* 17:555-572, 1999
57. Tsukamoto M, Inoue S, Ito T, et al: Functional evaluation of the bileaflet mechanical valve in the aortic position using dobutamine-stress echocardiography: Is a 23-mm prosthetic valve adequately large enough? *Jpn Circ J* 62:817-823, 1998
58. Cannon JD Jr, Zile MR, Crawford FA Jr, et al: Aortic valve resistance as an adjunct to the gorlin formula in assessing the severity of aortic stenosis in symptomatic patients. *J Am Coll Cardiol* 20:1517-1523, 1992
59. Burwash IG, Pearlman AS, Kraft CD, et al: Flow dependence of measures of aortic stenosis severity during exercise. *J Am Coll Cardiol* 24:1342-1350, 1994
60. de Filippi CR, Willett DL, Brickner ME, et al: Usefulness of dobutamine echocardiography in distinguishing severe from nonsevere valvular aortic stenosis in patients with depressed left ventricular function and low transvalvular gradients. *Am J Cardiol* 75:191-194, 1995
61. Lin SS, Roger VL, Pascoe R, et al: Dobutamine stress doppler hemodynamics in patients with aortic stenosis: Feasibility, safety, and surgical correlations [see comments]. *Am Heart J* 136:1010-1016, 1998
62. Leavitt JL, Coats MH, Falk RH: Effects of exercise on transmitral gradient and pulmonary artery pressure in patients with mitral stenosis or a prosthetic mitral valve: A doppler echocardiographic study. *J Am Coll Cardiol* 17:1520-1526, 1991
63. Tunick PA, Freedberg RS, Gargiulo A, et al: Exercise doppler echocardiography as an aid to clinical decision making in mitral valve disease. *J Am Soc Echocardiogr* 5:225-230, 1992
64. Dahan M, Paillole C, Martin D, et al: Determinants of stroke volume response to exercise in patients with mitral stenosis: A doppler echocardiographic study. *J Am Coll Cardiol* 21:384-389, 1993
65. Braverman AC, Thomas JD, Lee RT: Doppler echocardiographic estimation of mitral valve area during changing hemodynamic conditions. *Am J Cardiol* 68:1485-1490, 1991
66. Mangano DT: Perioperative cardiac morbidity. *Anesthesiology* 72:153-184, 1990
67. Mangano DT, London MJ, Tubau JF, et al: Dipyridamole thallium-201 scintigraphy as a preoperative screening test. A reexamination of its predictive potential. Study of perioperative ischemia research group [see comments]. *Circulation* 84:493-502, 1991
68. Marwick TH: Stress echocardiography with non-exercise techniques, in Otto CM (ed): *The Practice of Clinical Echocardiography*. Philadelphia, PA, Saunders, 1997, p 253
69. Das MK, Pellikka PA, Mahoney DW, et al: Assessment of cardiac risk before nonvascular surgery: Dobutamine stress echocardiography in 530 patients. *J Am Coll Cardiol* 35:1647-1653, 2000
70. Akosah KO, Olsovsky M, Kirshberg D, et al: Dobutamine stress echocardiography predicts cardiac events in heart transplant patients. *Circulation* 94:11283-11288, 1996
71. Derumeaux G, Redonnet M, Soyer R, et al: Assessment of the progression of cardiac allograft vasculopathy by dobutamine stress echocardiography. *J Heart Lung Transplant* 17:259-267, 1998
72. Spes CH, Klauss V, Mudra H, et al: Role of dobutamine stress echocardiography for diagnosis of cardiac allograft vasculopathy. *Transplant Proc* 30:904-906, 1998
73. Gunther F, Schwammenthal E, Rahmel A, et al: [initial experiences with dobutamine stress echocardiography in heart transplant patients] [Erste Erfahrungen mit der Dobutamin-Belastungsechokardiographie bei herztransplantierten Patienten.]. *Z Kardiol* 84:411-418, 1995
74. Verhoeven PP, Lee FA, Ramahi TM, et al: Prognostic value of noninvasive testing one year after orthotopic cardiac transplantation. *J Am Coll Cardiol* 28:183-189, 1996
75. Larsen RL, Applegate PM, Dyar DA, et al: Dobutamine stress echocardiography for assessing coronary artery disease after transplantation in children. *J Am Coll Cardiol* 32:515-520, 1998
76. Spes CH, Klauss V, Mudra H, et al: Diagnostic and prognostic value of serial dobutamine stress echocardiography for noninvasive assessment of cardiac allograft vasculopathy: A comparison with coronary angiography and intravascular ultrasound [see comments]. *Circulation* 100:509-515, 1999
77. Spes CH, Klauss V, Rieber J, et al: Functional and morphological findings in heart transplant recipients with a normal coronary angiogram: An analysis by dobutamine stress echocardiography, intracoronary doppler and intravascular ultrasound. *J Heart Lung Transplant* 18:391-398, 1999
78. Spes CH, Mudra H, Schnaack SD, et al: Dobutamine stress echocardiography for noninvasive diagnosis of cardiac allograft vasculopathy: A comparison with angiography and intravascular ultrasound. *Am J Cardiol* 78:168-174, 1996